



OSCE Checklist: Male Catheterisation

Introduction		
1	Introduce yourself to the patient including your name and role	
2	Confirm the patient's name and date of birth	
3	Briefly explain what the procedure will involve using patient-friendly language	
4	Explain the need for a chaperone	
5	Gain consent to proceed with catheterisation	
6	Check if the patient has any allergies	
7	Ask the patient if they have any pain before continuing with the clinical procedure	
8	Explain to the patient that they'll need to remove their underwear and lie on the clinical examination couch, covering themselves with the sheet provided	
9	Wash your hands using alcohol gel. If your hands are visibly soiled, wash them with soap and water	
Gather equipment		
10	Clean the top of a procedure trolley using an appropriate disinfectant wipe	
11	Collect the equipment required for the procedure and place it within reach on the clean trolley	
12	Check the expiry date on the catheter, sterile water, normal saline and lidocaine gel	
13	Ensure a clinical waste bin is placed nearby	
Equipment preparation		
14	Wash your hands again and don an apron	
15	Setup up the sterile field by first removing the outer packaging from the catheter pack and then opening the catheter pack from the corners without touching the inner surface of the field. Make sure to keep the catheter packaging as you'll need to transfer the sticky label containing the details of the catheter into the patient's notes	
16	Using aseptic non-touch technique (ANTT) empty the catheter, lidocaine gel syringe, sterile water syringe and sterile gloves onto the field	
17	Pour the 0.9% sodium chloride solution over the cotton balls which should already be located within the gallipot of the catheter pack	
Positioning the patient		
18	With the patient lying supine, ensure the bed is at an appropriate height for you to comfortably carry out the procedure	
19	Wash your hands again and don a pair of sterile gloves	
20	Ask your chaperone to remove the sheet covering the patient's genitals to allow you to maintain sterility	
21	Place a sterile absorbent pad underneath the patient's genital region, ensuring you maintain sterility	
Cleaning the penis		
22	Hold the penis with your non-dominant hand using some sterile gauze and ensure the patient's foreskin is retracted (if present)	

23	With your dominant hand pick up a cotton ball and use a single stroke moving away from the urethral meatus to clean an area of the glans. Dispose of the first cotton ball into the clinical waste bin and continue to repeat this process with a new cotton ball each time until all areas of the glans have been cleaned.	
24	Discard your used gloves, wash your hands again and don a new pair of sterile gloves	
25	Place the sterile drape over the patient's penis, positioned such that the penis remains visible through the central aperture of the drape. Some drapes come with a hole already present for this purpose, whereas others will require you to create one.	
26	Place the sterile urine collection bowl below the penis but on top of the sterile drape	
Inserting the anaesthetic		
27	Hold the penis vertically with your non-dominant hand using a piece of sterile gauze	
28	Warn the patient that the anaesthetic gel might initially sting, but then should quickly cause things to become numb	
29	With your dominant hand place the nozzle of the syringe of anaesthetic gel into the urethral meatus	
30	Empty the entire 10mls of anaesthetic gel into the urethra at a slow but steady pace	
31	Continue to hold to the penis in the vertical position to ensure the gel remains within the urethra and allow 3 to 5 minutes for the lidocaine gel to reach its maximum effect	
Inserting the catheter		
32	Pick up the catheter which should be on your sterile field in its wrapper	
33	Remove the tear-away portion of the wrapper near the catheter tip, making sure not to touch the catheter	
34	Hold the penis again using sterile gauze with your non-dominant 'dirty hand'	
35	Warn the patient you are about to insert the catheter	
36	Insert the exposed catheter tip into the urethral meatus using your dominant 'clean hand'	
37	Advance the catheter slowly whilst gradually removing more of the wrapper to expose more of the catheter	
38	You should continue to advance the catheter until it is fully inserted into the penis	
39	Once the catheter is fully inserted, inflate the catheter balloon with the 10ml syringe of sterile water to secure it within the bladder	
40	Once the balloon is fully inflated, remove the syringe and gently withdraw the catheter until resistance is noted, confirming the catheter is held securely within the bladder	
41	Attach the catheter bag tubing to the end of the catheter securely	
42	Position the catheter bag below the level of the patient to facilitate effective drainage of urine	
43	Replace the patient's retracted foreskin (if present) as failure to do so can result in the development of paraphimosis	
44	Clean away any urine spillage or excess lubricating gel and cover the patient with the sheet	
45	Dispose of your equipment into a clinical waste bin	
46	Wash your hands	
47	Provide the patient with privacy to get dressed	

To complete the procedure...

48	Explain to the patient that the procedure is now complete and that they should seek review if the catheter becomes painful or they feel unwell	
49	Thank the patient for their time	
50	Ask the nursing staff to monitor the patient's urine output and to contact you if they have any concerns	
51	Document the details of the procedure in the patient's notes and affix the sticky label from the catheter packaging beside your notes	

Read the full guide at
[geekymedics.com](https://www.geekymedics.com)



Download our
[clinical skills app](#)